

Names for H (ISBT 018) Blood Group Alleles

Intro

General description: The H blood group system consists of one antigen, H, that is carried on glycolipids and glycoproteins on the RBC membrane, where it is synthesised by the fucosyltransferase product of the *FUT1* gene; as well as on glycoproteins on epithelial cells and in body fluids, where it is synthesised by the fucosyltransferase product of the *FUT2* gene. In group O individuals, H antigen is the terminal antigen however, in group A and B individuals, the H antigen serves as the precursor structure for A and B blood-group-specific glycosyltransferases. Thus, group O people will test strongly H⁺ whereas groups A, B and AB will express very little H antigen. Mutations that negatively affect the α 2FucT1 enzyme activity (encoded by FUT1) will result in reduced or absent H production (and a concomitant decrease in A and/or B antigens in individuals where those enzymes are encoded). Total absence of H, A and B antigens is called the Oh or Bombay phenotype. Weak expression is referred to as the paraBombay phenotype. The enzymes α 2FucT1 and α 2FucT2 are single pass type II membrane glycoproteins in the Golgi. The α 2FucT1 protein consists of 365 amino acids and is encoded by FUT1 or H, if analysis is to predict a blood group antigen. The *FUT2* gene produces two transcripts; one of 343 amino acids and another more abundant form of 332 amino acids. The longer transcript encodes a protein with approximately one fourth the enzymatic activity and the shorter form is considered to be the active enzyme. However, in accordance with the LRG project it has been decided in 2017, we still keep to use the nucleotide for the longer transcript translation to count the position for each variant. The α 2FucT2 protein is encoded by FUT2 or Se, if analysis is to predict a blood group antigen.

Gene name: *FUT1*
 Number of exons: 4
 Initiation codon: Beginning of exon 4
 Stop codon: Within exon 4
 Entrez Gene ID: 2523
 LRG sequence: NG_007510.1 (genomic)
 NM_000148.3 (transcript)
 Reference allele: *FUT1*01* (shaded)
 Acceptable: *H* if inferred by hemagglutination/inhibition

Gene name: *FUT2*
 Number of exons: 2
 Initiation codon: Beginning of exon 2
 Stop codon: Within exon 2
 Entrez Gene ID: 2524

CAVE: In accordance with the LRG project it has been decided in 2017 that an ATG 33 nucleotides more 5' than the "old" ATG was the correct translation start ATG where the counting with 1 should be started. As a result, all *FUT2* sequences were to be corrected.

LRG sequence: NG_007511.1 (genomic)
 NM_000511.5 (transcript)
 Reference allele: *FUT2*01* (shaded)
 Acceptable: *Se* if inferred by hemagglutination/inhibition

Phenotype	Allele name	Nucleotide change	Exon Intron	Predicted amino acid change	(Reference No.) PMID	Accession number	rs number
H+	<i>FUT1*01</i>						
H+	<i>FUT1*01.02</i> (old: <i>FUT1*02</i>)	c.35C>T	4	p.Ala12Val	PMID: 15476160 PMID: 16331565		rs2071699
Weak phenotypes†							
H+weak	<i>FUT1*01W.01</i>	c.293C>T	4	p.Thr98Met	PMID: 15847661	AY611628	rs1469074926
H+weak	<i>FUT1*01W.02</i>	c.328G>A	4	p.Ala110Thr	PMID: 15476160 PMID: 34487549	DQ092446 DQ321370	rs56342683
H+weak	<i>FUT1*01W.03</i>	c.349C>T	4	p.His117Tyr	PMID: 9745152		rs150074056
H+weak	<i>FUT1*01W.04</i>	c.442G>T	4	p.Asp148Tyr	PMID: 9226185	AB004862	rs150074056
H+weak	<i>FUT1*01W.05.01</i>	c.460T>C	4	p.Tyr154His	PMID: 9031499		rs757349699
H+weak	<i>FUT1*01W.05.02</i>	c.460T>C c.1042G>A	4	p.Tyr154His p.Glu348Lys	PMID: 9226185 PMID: 9031498	AB004863	rs757349699 rs764739319
H+weak	<i>FUT1*01W.07</i>	c.491T>A	4	p.Leu164His	PMID: 7912436		rs104894687
H+weak	<i>FUT1*01W.08</i>	c.522C>A	4	p.Phe174Leu	PMID: 12366770	AF455028	rs747696745
H+weak	<i>FUT1*01W.09</i>	c.658C>T	4	p.Arg220Cys	PMID: 9031499		rs574691621
H+weak	<i>FUT1*01W.10</i>	c.659G>A	4	p.Arg220His	PMID: 11045762		rs1229284545
H+weak	<i>FUT1*01W.11</i>	c.661C>T	4	p.Arg221Cys	(5), Abstract		rs1452890889
H+weak	<i>FUT1*01W.12</i>	c.682A>G	4	p.Met228Val	PMID: 16403295 PMID: 17163878 PMID: 21988368		
H+weak	<i>FUT1*01W.13</i>	c.689A>C	4	p.Gln230Pro	PMID: 17176328		rs1445220556
H+weak	<i>FUT1*01W.14</i>	c.721T>C	4	p.Tyr241His	PMID: 9226185	AB004861	rs765114567
H+weak	<i>FUT1*01W.15</i>	c.801G>C	4	p.Trp267Cys	(1), Abstract (2), Abstract		
H+weak	<i>FUT1*01W.16</i>	c.801G>T	4	p.Trp267Cys	(1), Abstract (2), Abstract		

Phenotype	Allele name	Nucleotide change	Exon Intron	Predicted amino acid change	(Reference No.) PMID	Accession number	rs number
H+weak	<i>FUT1*01W.17</i>	c.832G>A	4	p.Asp278Asn	(1), Abstract (2), Abstract		
H+weak	<i>FUT1*01W.18</i>	c.903_904insAAC	4	p.Asn301_His302insAsn	n.a.	n.a.	
H+weak	<i>FUT1*01W.19</i>	c.917C>T	4	p.Thr306Ile	PMID: 17176328		
H+weak	<i>FUT1*01W.20</i>	c.990delG	4	p.Pro331Glnfs*6	PMID: 9226185	AB004860	
H+weak	<i>FUT1*01W.21</i>	c.235G>C	4	p.Gly79Arg	PMID: 20533259 PMID: 21988368	FJ665616	rs1399735219
H+weak	<i>FUT1*01W.22</i>	c.991C>A	4	p.Pro331Thr	PMID: 19572973	FM162557	rs1381389919
H+weak	<i>FUT1*01W.23</i>	c.424C>T	4	p.Arg142Trp	(3), Abstract PMID: 34487549	HQ891007	rs772921327
H+weak	<i>FUT1*01W.24</i>	c.649G>T	4	p.Val217Phe	PMID: 21839020 PMID: 34967725	GQ336988 HM584610	rs541722036
H+weak	<i>FUT1*01W.25</i>	obsolete					rs1399735219
H+weak	<i>FUT1*01W.26</i>	c.545G>A	4	p.Arg182His		KF385398	rs1284994775
H+weak	<i>FUT1*01W.27</i>	c.958G>A	4	p.Gly320Arg	PMID: 27893357	KF581194	rs762020231
H+weak	<i>FUT1*01W.28</i>	c.896A>C	4	p.Gln299Pro	PMID: 25858679 PMID: 30186784	KJ804401.1	
H+weak	<i>FUT1*01W.29</i>	c.655G>C	4	p.Val219Leu	PMID: 25538540	JX078970	
H+weak	<i>FUT1*01W.30</i>	avoided					
H+weak	<i>FUT1*01W.31</i> (old: <i>FUT1*02W.01</i>)	c.35C>T c.269G>T	4 4	p.Ala12Val p.Gly90Val	PMID: 17176328		rs897829842
H+weak	<i>FUT1*01W.32</i> (old: <i>FUT1*02W.02</i>)	c.35C>T c.371T>G	4 4	p.Ala12Val p.Phe124Cys	n.a.	n.a.	rs529462057
H+weak	<i>FUT1*01W.33</i> (old: <i>FUT1*02W.03</i>)	c.35C>T c.682A>G	4 4	p.Ala12Val p.Met228Val	PMID: 20533259		
H+weak	<i>FUT1*01W.34</i> (old: <i>FUT1*02W.05</i>)	c.35C>T c.748C>T	4 4	p.Ala12Val p.Arg250Trp	PMID: 25538540	JX317627	
H+weak	<i>FUT1*01W.35</i>	c.424C>T	4	p.Arg142Try	(6), Abstract		

Phenotype	Allele name	Nucleotide change	Exon Intron	Predicted amino acid change	(Reference No.) PMID	Accession number	rs number
H+weak	<i>FUT1*01W.36</i>	c.755G>C	4	p.Arg253Pro	PMID: 30217757		
H+weak	<i>FUT1*01W.37</i>	c.803G>A	4	p.Cys268Tyr	PMID: 34693534	MW821348	
H+weak	<i>FUT1*01W.38</i>	c.814A>G	4	p.Ile272Val	PMID: 30217757		
H+weak	<i>FUT1*01W.39</i>	c.229C>T c.302C>T	4	p.Leu77Phe Ala101Val	PMID: 34792200	MN971576	
Null alleles							
H-	<i>FUT1*01N.01</i>	c.422G>A	4	p.Trp141Ter	PMID: 17176328		rs749165173
H-	<i>FUT1*01N.02</i>	c.461A>G	4	p.Tyr154Cys	PMID: 9122901		
H-	<i>FUT1*01N.03</i>	c.462C>A	4	p.Tyr154Ter	(4), Abstract		
H-	<i>FUT1*01N.04</i>	c.513G>C	4	p.Trp171Cys	PMID: 9122901		
H-	<i>FUT1*01N.05</i>	c.538C>T	4	p.Gln180Ter	PMID: 17176328		rs746629771
H- /H+w (‡)	<i>FUT1*01N.06</i>	c.551_552delAG	4	p.Glu184Valfs*85	PMID: 9031499		rs573412368
H-	<i>FUT1*01N.07</i>	c.586C>T	4	p.Gln196Ter	PMID: 15476160	DQ157279	
H-	<i>FUT1*01N.08</i>	c.695G>A	4	p.Trp232Ter	PMID: 9226185	AB004859	
H-	<i>FUT1*01N.09</i>	c.725T>G	4	p.Leu242Arg	PMID: 9299444 PMID: 9745152		rs28934588
H-	<i>FUT1*01N.10</i>	c.776T>A	4	p.Val259Glu	PMID: 9122901		
H-	<i>FUT1*01N.11</i>	c.785G>A c.786C>A	4	p.Ser262Lys	PMID: 11161242	AJ276886	
H-	<i>FUT1*01N.12</i>	c.826C>T	4	p.Gln276Ter	PMID: 7912436		rs104894688
H-	<i>FUT1*01N.13</i>	c.881_882delTT	4	p.Phe294Cysfs*40	PMID: 9031499		rs777455020
H-	<i>FUT1*01N.14</i>	c.944C>T	4	p.Ala315Val	PMID: 9122901		
H-	<i>FUT1*01N.15</i>	c.948C>G	4	p.Tyr316Ter	PMID: 7912436		rs104894686

Phenotype	Allele name	Nucleotide change	Exon Intron	Predicted amino acid change	(Reference No.) PMID	Accession number	rs number
H-	FUT1*01N.16 (old: FUT1*02W.04)	c.980A>C	4	p.Asn327Thr	PMID: 15487706 PMID: 23560544 PMID: 9031499		rs777291875
H-	FUT1*01N.17	c.1047G>C	4	p.Trp349Cys	PMID: 9122901		rs1438752561
H-	FUT1*01N.18	c.684G>A	4	p.Met228Ile	n.a.		
H-	FUT1*01N.19	c.694T>C	4	p.Trp232Arg	n.a.		
H-	FUT1*01N.20	c.768delC	4	p.Val257Phefs*23	PMID: 34967725	KM514482	
H-	FUT1*01N.21	c.13_19dup	4	p.Arg7Glnfs*63	n.a.		rs150995632
H-	FUT1*01N.22	c.791_792insG	4	p.Met265Hisfs*5		MG987419	
H-	FUT1*01N.23	c.710delG	4	p.Gly237Alafs*43		MH298872	rs1486913817
H-	FUT1*01N.24	c.454delG	4	p.Glu152Argfs*6		MH298873	rs1409390706
H-	FUT1*01N.25	c.288T>A	4	p.Tyr96Ter		MH298874	
H-	FUT1*01N.26 (old: FUT1*02N.01)	c.35C>T c.423G>A	4 4	p.Ala12Val p.Trp141Ter	PMID: 21839020	HQ699894	rs2071699
H-	FUT1*01N.27	c.49T>C	4	p.Val17Arg	PMID: 29441582	KX644898	
H-	FUT1*01N.28	c.361G>A	4	p.Ala121Thr	PMID: 34539321	MN938362	
H-	FUT1*01N.29	c.366-398del33	4	p.Val123Glu fsX355	PMID:17922418		
H-	FUT1*01N.30	c.392T>C	4	p.Leu131Pro	PMID: 26926997		
H-	FUT1*01N.31	c.508dupT	4	p. 170 fsX268	PMID: 33175455	KM255205	
H-	FUT1*01N.32	c.668-670delACT	4	p.Tyr224del	PMID: 26926997 PMID: 30988570		
H-	FUT1*01N.33	c.749-765delGGGC ACGGCACGAAGCC	4		PMID: 34799859	MT078122	
H-	FUT1*01N.34	c.787A>C	4	p.Asn263His	PMID: 33175455	KM255206	
H-	FUT1*01N.35	c.985insG c.961G>A	4 4	p.319Glyfs334 p.Asp321Asn	PMID: 28026021	KT989645	

Phenotype	Allele name	Nucleotide change	Exon Intron	Predicted amino acid change	(Reference No.) PMID	Accession number	rs number
H-	<i>FUT1*01N.36</i>	c.1051G>T	4	p.Gly351Cys	PMID: 30988570		
H-	<i>FUT1*01N.37</i>	c.35C>T c.980A>C	4 4	p.Ala12Val p.Asn327Thr	PMID: 15487706 PMID: 23560544 PMID: 9031499		rs777291875
<p>‡ PMID: 32110200: Adsorption-elution studies with the patient's RBCs revealed positive reactions with anti-B, confirming a weak B phenotype. Flow cytometric analysis using fluorochrome-conjugated anti-H (BRIC198) detected minimal amounts of H antigens on the patient's RBCs with homozygous c.515_516delAG of FUT1.</p>							
Null phenotypes – Gene deletions							
H-	<i>FUT1*0N.01</i>	Coding region deletion (5443-bp length of deletion including the 1098-bp coding sequence of FUT1 gene)	-	p.0	PMID: 26926997		
<p>† Note that H expression will be masked if a functional A or B allele is also inherited. Also, that H antigen may be weakly detectable on RBCs where FUT*01N homozygosity occurs, due to the adsorption of soluble H antigen synthesized by FUT2.</p>							
Phenotype (saliva) †	Allele name	Nucleotide change	Exon Intron	Predicted amino acid change	(Reference No.) PMID	Accession number	rs number
H+	<i>FUT2*01</i>						
H+	<i>FUT2*01.02</i> (old: <i>FUT2*02</i>)	c.37G>A	2	p.Ala13Thr	PMID: 18422843		rs532291838
H+	<i>FUT2*01.03.01</i> (old: <i>FUT2*03.01</i>)	c.73A>G	2	p.Ile25Val	PMID: 9760207		rs1800021
H+	<i>FUT2*01.03.02</i> (old: <i>FUT2*03.02</i>)	c.73A>G c.146C>T	2	p.Ile25Val p.Ala49Val	PMID: 9760207		rs1800021 rs114018037
H+	<i>FUT2*01.03.03</i> (old: <i>FUT2*03.03</i>)	c.73A>G c.514G>A	2	p.Ile25Val p.Asp172Asn			rs1800021 rs1800025
H+	<i>FUT2*01.04</i> (old: <i>FUT2*04</i>)	c.412C>T	2	p.Arg138Cys	PMID: 9760207		rs1800022

Phenotype	Allele name	Nucleotide change	Exon Intron	Predicted amino acid change	(Reference No.) PMID	Accession number	rs number
H+	<i>FUT2*01.05</i> (old: <i>FUT2*05</i>)	c.433G>A	2	p.Val145Ile	PMID: 10980544 PMID: 1456946		rs370886251
H+	<i>FUT2*01.06</i> (old: <i>FUT2*06</i>)	c.514G>A	2	p.Asp172Asn	PMID: 9760207		rs1800025
H+	<i>FUT2*01.07</i> (old: <i>FUT2*07</i>)	c.698G>A	2	p.Arg233His	PMID: 18422843		rs776245547
H+	<i>FUT2*01.08</i> (old: <i>FUT2*08</i>)	c.718G>A	2	p.Val240Met	PMID: 10980544		rs375360260
H+	<i>FUT2*01.09</i> (old: <i>FUT2*09</i>)	c.749G>A	2	p.Arg250Gln	PMID: 15476160	DQ321371	rs369911091
H+	<i>FUT2*01.10</i> (old: <i>FUT2*10</i>)	c.780_781nsGTG	2	p.260_261insVal	PMID: 17655580		
H+	<i>FUT2*01.11</i>	c.131A>G c.134T>G	2 2	p.Gln44Arg p.Ile45Arg	PMID: 34487549	MW309872	
Weak phenotypes							
H+w	<i>FUT2*01W.01</i>	c.311C>T	2	p.Ala104Val			rs149356814
H+w	<i>FUT2*01W.02.01</i>	c.418A>T	2	p.Ile140Phe			rs1047781
H+w	<i>FUT2*01W.02.02</i>	c.418A>T c.650T>G	2 2	p.Ile140Phe p.Val217Gly			rs1047781
H+w	<i>FUT2*01W.02.03</i>	c.418A>T c.874G>A	2 2	p.Ile140Phe p.Gly292Arg			rs1047781 rs548111243
H+w	<i>FUT2*01W.03</i>	c.886G>A	2	p.Ala296Thr			rs79097987
H+w	<i>FUT2*01W.04</i>	c.650T>C	4	p.Val217Ala	PMID: 34487549	MW661069	
Null phenotypes – Nucleotide polymorphisms							
H-	<i>FUT2*01N.01</i>	c.277G>A c.418A>T	2	p.Ala93Thr p.Ile140Phe			rs112722916 rs1047781
H-	<i>FUT2*01N.02</i>	c.461G>A	2	p.Trp154Ter	PMID: 7876235	U17894	rs601338

Phenotype	Allele name	Nucleotide change	Exon Intron	Predicted amino acid change	(Reference No.) PMID	Accession number	rs number
H-	<i>FUT2*01N.03</i>	c.602G>A	2	p.Arg201His	PMID: 18422843		rs572832908
H-	<i>FUT2*01N.04</i>	c.604C>T	2	p.Arg202Ter	PMID: 8755920 PMID: 10550557 PMID: 8928486 PMID: 8670215 PMID: 11606829		rs1800028
H-	<i>FUT2*01N.05</i>	c.661C>T	2	p.Arg221Ter	PMID: 8755920		rs1800029
H-	<i>FUT2*01N.06</i>	c.691C>T	2	p.Arg231Ter	PMID: 10319583 PMID: 34487549		rs144566043
H-	<i>FUT2*01N.07</i>	c.697C>T	2	p.Arg233Cys	PMID: 14569463		rs768236330
H-	<i>FUT2*01N.08</i>	c.718_719delGT	2	p.Val240Glyfs*4	PMID: 10085528 PMID: 1160682		
H-	<i>FUT2*01N.09</i>	c.721_723delGTC	2	p.Val241del	PMID: 10550557		
H-	<i>FUT2*01N.10</i>	c.433G>A c.793G>A	2 2	p.Val145Ile p.Asp265Asn	PMID: 14569463		rs370886251 rs907232085
H-	<i>FUT2*01N.11</i>	c.811delC	2	p.Pro271Leufs*16	PMID: 9760207		
H-	<i>FUT2*01N.12</i>	c.882G>A	2	p.Trp294Ter	PMID: 10550557 PMID: 8670215 PMID: 11606829		rs1800030
H-	<i>FUT2*01N.13</i>	c.901G>A	2	p.Gly301Arg	PMID: 14569463		rs144269088
H-	<i>FUT2*01N.14</i>	c.983C>T	2	p.Pro328Leu	PMID: 18422843		rs200626231
H-	<i>FUT2*01N.15</i>	c.335C>T	2	p.Pro112Leu	PMID: 10533829		rs200157007
H-	<i>FUT2*01N.16</i>	c.772G>A	2	p.Gly258Ser	PMID: 20570966		rs60266286
H-	<i>FUT2*01N.17</i>	c.445G>A	2	p.Gly149Ser	PMID: 19572973	FM180558	rs200543547
H-	<i>FUT2*01N.18</i>	c.851C>A	2	p.Thr284Asn	PMID: 22188519		rs371279676

Phenotype	Allele name	Nucleotide change	Exon Intron	Predicted amino acid change	(Reference No.) PMID	Accession number	rs number
Null phenotypes – Gene deletions							
H–	<i>FUT2*0N.01</i>	Gene deletion		p.0			
H–	<i>FUT2*0N.01</i>	Coding region deleted		p.0	PMID: 9299444 PMID: 9745152 PMID: 10982186		
H–	<i>FUT2*0N.03</i>	Fusion gene 1 between <i>FUT2</i> and <i>Sec1</i>		-	PMID: 8755920 PMID: 18067503	D82933	
H–	<i>FUT2*0N.04</i>	Fusion gene 2 between <i>FUT2</i> and <i>Sec1</i>		-			
H–	<i>FUT2*0N.05</i>	Deletion of <i>FUT2</i>		p.0	PMID: 26926997		

† Saliva phenotype is shown here to represent secreted H antigen in all body fluids

References

- Abstract (1) Johnson PH & et al. *Vox Sang* 1994;67(Suppl 2):25.
- Abstract (2) cited according to Scharberg et al. *Immunohematology* 2016; 32(3):112-118. The abstract itself contains insufficient information.
- Abstract (3) Chi Q & et al. *Vox Sang* 2011;101(Suppl 2):108.
- Abstract (4) Ogasawara K & et al. *Vox Sang* 2000;78(Suppl 1):4.
- Abstract (5) Samuelsson J & et al. Annual meeting of Australian and New Zealand Society for Blood Transfusion, Sydney, Australia, October 2005: (former) Abstract 47.
- Abstract (6) Jin S, Liu X, Xiang D. Identification of a novel and rare FUT1 allele in one Chinese Han Para-Bombay pedigree. *Chin J Blood Transfus*, 2016; 29(Suppl 1):110
- PMID 1160682 G H Chye, S T Hong. Salmonella typhi meningitis. A case report and family investigations. *Med J Malaysia*. 1975 Mar;30(3):219-22.
- PMID 1456946 J M Argilés, N Carbó, F J López-Soriano. Hepatic accumulation of alpha-aminoisobutyrate in the pregnant rat at term. *Biochem Int*. 1992 Oct;28(2):199-204.
- PMID 7876235 R J Kelly, S Rouquier, D Giorgi & et al. Sequence and expression of a candidate for the human Secretor blood group alpha(1,2)fucosyltransferase gene (FUT2). Homozygosity for an enzyme-inactivating nonsense mutation commonly correlates with the non-secretor phenotype. *J Biol Chem*. 1995 Mar 3;270(9):4640-9. doi: 10.1074/jbc.270.9.4640.
- PMID 7912436 R J Kelly, L K Ernst, R D Larsen & et al. Molecular basis for H blood group deficiency in Bombay (Oh) and para-Bombay individuals. *Proc Natl Acad Sci U S A*. 1994 Jun 21;91(13):5843-7. doi: 10.1073/pnas.91.13.5843.
- PMID 8670215 L C Yu, R E Broadberry, Y H Yang & et al. Heterogeneity of the human Secretor alpha(1,2)fucosyltransferase gene among Lewis(a+b-) non-secretors. *Biochem Biophys Res Commun*. 1996 May 15;222(2):390-4. doi: 10.1006/bbrc.1996.0754.
- PMID 8755920 Y Koda, M Soejima, Y Liu, H Kimura. Molecular basis for secretor type alpha(1,2)-fucosyltransferase gene deficiency in a Japanese population: a fusion gene generated by unequal crossover responsible for the enzyme deficiency. *Am J Hum Genet*. 1996 Aug;59(2):343-50.
- PMID 8928486 S Henry, R Mollicone, J B Lowe & et al. A second nonsecretor allele of the blood group alpha(1,2)fucosyl-transferase gene (FUT2). *Vox Sang*. 1996;70(1):21-5. doi: 10.1111/j.1423-0410.1996.tb00991.x.
- PMID 9031498 B Wang, Y Koda, M Soejima, H Kimura. Two missense mutations of H type alpha(1,2)fucosyltransferase gene (FUT1) responsible for para-Bombay phenotype. *Vox Sang*. 1997;72(1):31-5. doi: 10.1046/j.1423-0410.1997.00031.x.

- PMID 9031499 L C Yu, Y H Yang, R E Broadberry & et al. Heterogeneity of the human H blood group alpha(1,2)fucosyltransferase gene among para-Bombay individuals. *Vox Sang.* 1997;72(1):36-40. doi: 10.1046/j.1423-0410.1997.00036.x.
- PMID 9122901 F F Wagner, W A Flegel. Polymorphism of the h allele and the population frequency of sporadic nonfunctional alleles. *Transfusion.* 1997 Mar;37(3):284-90. doi: 10.1046/j.1537-2995.1997.37397240210.x.
- PMID 9226185 M Kaneko, S Nishihara, N Shinya, T Kudo, H Iwasaki, T Seno, Y Okubo, H Narimatsu. Wide variety of point mutations in the H gene of Bombay and para-Bombay individuals that inactivate H enzyme. *Blood.* 1997 Jul 15;90(2):839-49.
- PMID 9299444 Y Koda, M Soejima, P H Johnson & et al. Missense mutation of FUT1 and deletion of FUT2 are responsible for Indian Bombay phenotype of ABO blood group system. *Biochem Biophys Res Commun.* 1997 Sep 8;238(1):21-5. doi: 10.1006/bbrc.1997.7232.
- PMID 9745152 P Fernandez-Mateos, A Cailleau, S Henry & et al. Point mutations and deletion responsible for the Bombay H null and the Reunion H weak blood groups. *Vox Sang.* 1998;75(1):37-46.
- PMID 9760207 Y Liu, Y Koda, M Soejima & et al. Extensive polymorphism of the FUT2 gene in an African (Xhosa) population of South Africa. *Hum Genet.* 1998 Aug;103(2):204-10. doi: 10.1007/s004390050808.
- PMID 10085528 L C Yu, H L Lee, C C Chu & et al. A newly identified nonsecretor allele of the human histo-blood group alpha(1,2)fucosyltransferase gene (FUT2). *Vox Sang.* 1999;76(2):115-9. doi: 10.1159/000031030.
- PMID 10319583 Y H Liu, Y Koda, M Soejima & et al. The fusion gene at the ABO-secretor locus (FUT2): absence in Chinese populations. *J Hum Genet.* 1999;44(3):181-4. doi: 10.1007/s100380050138.
- PMID 10533829 J G Chang, T Y Yang, T C Liu & et al. Molecular analysis of secretor type alpha(1,2)-fucosyltransferase gene mutations in the Chinese and Thai populations. *Transfusion.* 1999 Sep;39(9):1013-7. doi: 10.1046/j.1537-2995.1999.39091013.x.
- PMID 10550557 C T Peng, C H Tsai, T P Lin & et al. Molecular characterization of secretor type alpha(1, 2)-fucosyltransferase gene deficiency in the Philippine population. *Ann Hematol.* 1999 Oct;78(10):463-7. doi: 10.1007/s002770050599.
- PMID 10980544 H Pang, N Fujitani, M Soejima & et al. Two distinct Alu-mediated deletions of the human ABO-secretor (FUT2) locus in Samoan and Bangladeshi populations. *Hum Mutat.* 2000 Sep;16(3):274. doi: 10.1002/1098-1004(200009)16:3<274::AID-HUMU20>3.0.CO;2-I.
- PMID 10982186 Y Koda, M Soejima, P H Johnson & et al. An Alu-mediated large deletion of the FUT2 gene in individuals with the ABO-Bombay phenotype. *Hum Genet.* 2000 Jan;106(1):80-5. doi: 10.1007/s004390051013.
- PMID 11045762 C F Sun, M D Lo, C H Lee, D C Chu. Novel mutations, including a novel G659A missense mutation, of the FUT1 gene are responsible for the para-Bombay phenotype. *Ann Clin Lab Sci.* 2000 Oct;30(4):387-90.

- PMID 11161242 T Wagner, M Vadon, E Staudacher, A Schmarda, C Gassner, W Helmberg, G Lanzer, W A Flegel, F F Wagner. A new h allele detected in Europe has a missense mutation in alpha(1,2)-fucosyltransferase motif II. *Transfusion*. 2001 Jan;41(1):31-8. doi: 10.1046/j.1537-2995.2001.41010031.x.
- PMID 11606829 L C Yu, C C Chu, Y S Chan & et al. Polymorphism and distribution of the Secretor alpha(1,2)-fucosyltransferase gene in various Taiwanese populations. *Transfusion*. 2001 Oct;41(10):1279-84. doi: 10.1046/j.1537-2995.2001.41101279.x.
- PMID 12366770 S P Yip, K Y Chee, P Y Chan, E Y D Chow, H F Wong. Molecular genetic analysis of para-Bombay phenotypes in Chinese: a novel non-functional FUT1 allele is identified. *Vox Sang*. 2002 Oct;83(3):258-62. doi: 10.1046/j.1423-0410.2002.00184.x.
- PMID 14569463 Yoshiro Koda, Takafumi Ishida, Hidenori Tachida & et al. DNA sequence variation of the human ABO-secretor locus (FUT2) in New Guinean populations: possible early human migration from Africa. *Hum Genet*. 2003 Nov;113(6):534-41. doi: 10.1007/s00439-003-1013-6. Epub 2003 Sep 3.
- PMID 15476160 Zhong-hui Guo, Dong Xiang, Zi-yan Zhu & et al. [Analysis on FUT1 and FUT2 gene of 10 para-Bombay individuals in China]. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi*. 2004 Oct;21(5):417-21.
- PMID 15487706 Ding-Ping Chen, Ching-Ping Tseng, Wei-Ting Wang & et al. Two prevalent h alleles in para-Bombay haplotypes among 250,000 Taiwanese. *Ann Clin Lab Sci*. Summer 2004;34(3):314-8.
- PMID 15847661 Lixing Yan, Faming Zhu, Xianguo Xu, Xiaozhen Hong, Qinfeng Lv. Molecular basis for para-Bombay phenotypes in Chinese persons, including a novel nonfunctional FUT1 allele. *Transfusion*. 2005 May;45(5):725-30. doi: 10.1111/j.1537-2995.2005.04305.x.
- PMID 16331565 Xian-guo Xu, Xiao-zhen Hong, Jun-jie Wu, Kai-rong Ma, Qi-hua Fu, Li-xing Yan. [Study on the frequency of alpha-1,2-fucosyltransferase gene h4 allele (C35T) in Chinese population]. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi*. 2005 Dec;22(6):657-60.
- PMID 16403295 Xiao-Zhen Hong, Xiao-Chun Shao, Xian-Guo Xu & et al. [Study on the molecular genetics basis for one para-Bombay phenotype]. *Zhongguo Shi Yan Xue Ye Xue Za Zhi*. 2005 Dec;13(6):1120-4.
- PMID 17176328 Jill R Storry, Jannica S Johannesson, Joyce Poole & et al. Identification of six new alleles at the FUT1 and FUT2 loci in ethnically diverse individuals with Bombay and Para-Bombay phenotypes. *Transfusion*. 2006 Dec;46(12):2149-55. doi: 10.1111/j.1537-2995.2006.01045.x.
- PMID 17655580 Shea Ping Yip, Siu Kin Lai, Maria L Wong. Systematic sequence analysis of the human fucosyltransferase 2 (FUT2) gene identifies novel sequence variations and alleles. *Transfusion*. 2007 Aug;47(8):1369-80. doi: 10.1111/j.1537-2995.2007.01280.x.
- PMID 18067503 Mikiko Soejima, Junko Fujihara, Haruo Takeshita, Yoshiro Koda. Sec1-FUT2-Sec1 hybrid allele generated by interlocus gene conversion. *Transfusion*. 2008 Mar;48(3):488-92. doi: 10.1111/j.1537-2995.2007.01553.x. Epub 2007 Dec 7.
- PMID 18422843 Mikiko Soejima, Tamiko Nakajima, Junko Fujihara & et al. Genetic variation of FUT2 in Ovambos, Turks, and Mongolians. *Transfusion*. 2008 Jul;48(7):1423-31. doi: 10.1111/j.1537-2995.2008.01710.x. Epub 2008 Apr 18.

- PMID 19572973 Eva Maria Matzhold, Wolfgang Helmberg, Thomas Wagner & et al. Identification of 14 new alleles at the fucosyltransferase 1, 2, and 3 loci in Styrian blood donors, Austria. *Transfusion*. 2009 Oct;49(10):2097-108. doi: 10.1111/i.1537-2995.2009.02293.x. Epub 2009 Jun 30.
- PMID 20533259 Xian-guo Xu, Xiao-zhen Hong, Ying Liu & et al. [Analysis of alpha-1,2-fucosyltransferase gene mutations in a Chinese family with para-Bombay phenotype]. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi*. 2010 Jun;27(3):250-4. doi: 10.3760/cma.j.issn.1003-9406.2010.0.003.
- PMID 21839020 Xiao-Hong Cai, Sha Jin, Xi Liu & et al. Molecular genetic analysis for the para-Bombay blood group revealing two novel alleles in the FUT1 gene. *Blood Transfus*. 2011 Oct;9(4):466-8. doi: 10.2450/2011.0115-10. Epub 2011 Jun 15.
- PMID 22188519 Mikiko Soejima, Ryo Fujimoto, Tetsuro Agusa & et al. Genetic variation of FUT2 in a Vietnamese population: identification of two novel Se enzyme-inactivating mutations. *Transfusion*. 2012 Jun;52(6):1268-75. doi: 10.1111/j.1537-2995.2011.03485.x. Epub 2011 Dec 21.
- PMID 23560544 Guangping Luo, Ling Wei, Zhen Wang & et al. The summary of FUT1 and FUT2 genotyping analysis in Chinese para-Bombay individuals including additional nine probands from Guangzhou in China. *Transfusion*. 2013 Dec;53(12):3224-9. doi: 10.1111/trf.12183. Epub 2013 Apr 8.
- PMID 25538540 Ziyi He, Fuping Liu. Two Novel α 1,2-Fucosyltransferase Alleles in an H-Deficient Phenotype Individual. *Transfus Med Hemother*. 2014 Oct;41(5):375-9. doi: 10.1159/000366235. Epub 2014 Aug 17.
- PMID 25858679 Wei Liang, Hong Xu, Yi-yu Liu, Shou-rong Zhou & et al. Molecular genetic analysis of para-Bombay phenotype in Chinese persons: a novel FUT1 allele is identified. *Transfusion*. 2015 Jun;55(6 Pt 2):1588. doi: 10.1111/trf.13022. Epub 2015 Apr 9
- PMID 27893357 Bogumila Michalewska, Martin L Olsson, Grazyna Naremska & et al. FUT1 mutations responsible for the H-deficient phenotype in the Polish population, including the first example of an abolished start codon. *Blood Transfus*. 2018 Jan;16(1):101-104. doi: 10.2450/2016.0135-16. Epub 2016 Nov 21.
- PMID 34792200 Qing Chen, Chenchen Feng, Kshitij Srivastava & et al. NM_000148.4(FUT1):c.[229C>T;302C>T] with 2 missense variations in the FUT1 gene associated with a para-Bombay phenotype. *Transfusion*. 2022 Jan;62(1):E5-E7. doi: 10.1111/trf.16743. Epub 2021 Nov 18.
- PMID 17163878 Yan LX, Xu XG, Hong XZ, Wu JJ, Zhu FM, Fu QH. Identification of a novel FUT1 allele derived from the alpha-(1,2)-fucosyltransferase gene through a nucleotide substitution 682A>G. *Transfus Med*. 2006 Dec;16(6):447-9. doi: 10.1111/j.1365-3148.2006.00706.x. PMID: 17163878.
- PMID 21988368 Xu X, Tao S, Ying Y, Hong X, He Y, Zhu F, Lv H, Yan L. A novel FUT1 allele was identified in a Chinese individual with para-Bombay phenotype. *Transfus Med*. 2011 Dec;21(6):385-93. doi: 10.1111/j.1365-3148.2011.01111.x. Epub 2011 Oct 12. PMID: 21988368.
- PMID 34967725 Sun X, Cai Y, Ni H, Cao X, Ma Y, Zhang Y, Jiang C, Lu J, Cong H. Analysis of the molecular mechanism and pedigree investigation of para-Bombay phenotype caused by combined mutations at position h649 and h768 of FUT1 gene. *Blood Transfus*. 2021 Nov 29. doi: 10.2450/2021.0215-21. Epub ahead of print. PMID: 34967725.

- PMID 25858679 Liang W, Xu H, Liu YY, Zhou SR, Deng DF, Yang L, Xu DY, Deng G, Mei CL, He YL, Zhang Z. Molecular genetic analysis of para-Bombay phenotype in Chinese persons: a novel FUT1 allele is identified. *Transfusion*. 2015 Jun;55(6 Pt 2):1588. doi: 10.1111/trf.13022. Epub 2015 Apr 9. PMID: 25858679.
- PMID 30186784 Liang W, Cai F, Yang L, Zhang Z, Wang Z. Four Non-functional FUT1 Alleles Were Identified in Seven Chinese Individuals with Para-Bombay Phenotypes. *Iran J Public Health*. 2018 Aug;47(8):1128-1136. PMID: 30186784; PMCID: PMC6123583.
- PMID 30217757 Lin J, Chen R, Zhu S, Qiu X, Huang Y, Wang X. Sequence analysis of α -(1, 2)-fucosyltransferase gene in nine Chinese individuals with Para-Bombay phenotype. *Gene*. 2019 Feb 15;685:21-23. doi: 10.1016/j.gene.2018.09.017. Epub 2018 Sep 12. PMID: 30217757.
- PMID 34693534 Han B, Zhu J, Zhu Y, Feng Z, Zhang S. A novel FUT1 variant allele was identified in a Chinese Para-Bombay individual. *Transfusion*. 2022 Jan;62(1):E1-E2. doi: 10.1111/trf.16712. Epub 2021 Oct 24. PMID: 34693534.
- PMID 29441582 Hong X, Chen S, Ma K, He J, Zhu F. c.49T>C mutation on the α -(1,2)-fucosyltransferase gene responsible for an individual with para-Bombay phenotype. *Transfusion*. 2018 Apr;58(4):1076-1077. doi: 10.1111/trf.14499. Epub 2018 Feb 13. PMID: 29441582.
- PMID 34539321 Lei H, Shen Y, Wang Y, Su N, Wang X, Cai X. A Para-Bombay Blood Group Case Associated with a Novel FUT1 Mutation c.361G>A. *Transfus Med Hemother*. 2021 Aug;48(4):254-258. doi: 10.1159/000513318. Epub 2021 Jan 13. PMID: 34539321; PMCID: PMC8406362.
- PMID 17922418 Su Yq, Wei TL, Yu Q, Liang YL, Li DC. [Molecular genetic analysis of FUT1 and FUT2 gene in para-Bombay Chinese: a novel FUT1 allele is identified]. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi*. 2007 Oct;24(5):520-3. Chinese. PMID: 17922418.
- PMID 26926997 Zanjani DS, Afzal Aghaee M, Badiie Z, Mehrasa R, Roodsarabi A, Khayyami ME, Shahabi M. Molecular basis of Bombay phenotype in Mashhad, Iran: identification of a novel FUT1 deletion. *Vox Sang*. 2016 Jul;111(1):88-92. doi: 10.1111/vox.12393. Epub 2016 Feb 29. PMID: 26926997.
- PMID 33175455 Lin FQ, Zhang KL, Li XF, Zhang X, Liu Y, Li JP. Two novel FUT1 alleles that cause para-Bombay phenotype in a Chinese individual. *Transfusion*. 2020 Dec;60(12):E55-E57. doi: 10.1111/trf.16183. Epub 2020 Nov 11. PMID: 33175455.
- PMID 34799859 Hu B, Yin F, Liu L, Feng Z, Pang S. Identification of a novel FUT1 allele in a Chinese individual with para-Bombay phenotype. *Transfusion*. 2022 Mar;62(3):E19-E20. doi: 10.1111/trf.16750. Epub 2021 Nov 19. PMID: 34799859.
- PMID 28026021 Wang L, Huang C, Wei Z, Tan J, Qin L, Tian L. Identification of a novel FUT1 allele with two mutations in a Chinese para-Bombay individual. *Transfusion*. 2017 Jan;57(1):93-96. doi: 10.1111/trf.13879. Epub 2016 Dec 27. PMID: 28026021.
- PMID 30988570 Roshan Mehr F, Manafi M, Sharifi Z, Shahabi M. Mutational Analysis of Bombay Phenotype in Iranian People: Identification of a Novel FUT1 Allele. *Indian J Hematol Blood Transfus*. 2019 Apr;35(2):321-324. doi: 10.1007/s12288-018-1027-9. Epub 2018 Oct 8. PMID: 30988570; PMCID: PMC6439033.

- PMID 34487549 Anukul N, Wita R, Leetrakool N, Sirikul C, Veeraphan N, Wongchai S. Two novel alleles on Fucosyltransferase 2 from northern Thai para-Bombay family and computational prediction on mutation effect. *Transfusion*. 2021 Nov;61(11):3247-3257. doi: 10.1111/trf.16646. Epub 2021 Sep 6. PMID: 34487549.
- PMID 20570966 McGovern DP, et al. Fucosyltransferase 2 (FUT2) non-secretor status is associated with Crohn's disease. *Hum Mol Genet*. 2010 Sep 1;19(17):3468-76.

Track of changes		from	to
1	Version	v6.0 30-JUN-2022	v6.1 31-MAR-2023
2	Author	created Franz Wagner, Yanli Ji, June 2022	Yanli Ji, March 2023
3	Reviewer	reviewed Franz Wagner, C. Gassner, June 2022	Franz Wagner, March 2023
4	Intro	changed	CAVE noted
5	Allele	corrected <i>FUT2*01.02</i> , c.4G>A, p.Ala2Thr	<i>FUT2*01.02</i> , c.37G>A, p.Ala13Thr
6	Allele	corrected <i>FUT2*01.03.01</i> , c.40A>G, p.Ile14Val	<i>FUT2*01.03.01</i> , c.73A>G, p.Ile25Val
7	Allele	corrected <i>FUT2*01.03.02</i> , c.40A>G, c.113C>T, p.Ile14Val, p.Ala38Val	<i>FUT2*01.03.02</i> , c.73A>G, c.146C>T, p.Ile25Val, p.Ala49Val
8	Allele	corrected <i>FUT2*01.03.03</i> , c.40A>G, c.481G>A, p.Ile14Val, p.Asp161Asn	<i>FUT2*01.03.03</i> , c.73A>G, c.514G>A, p.Ile25Val, p.Asp172Asn
9	Allele	corrected <i>FUT2*01.04</i> , c.379C>T, p.Arg127Cys	<i>FUT2*01.04</i> , c.412C>T, p.Arg138Cys
10	Allele	corrected <i>FUT2*01.05</i> , c.400G>A, p.Val134Ile	<i>FUT2*01.05</i> , c.433G>A, p.Val145Ile
11	Allele	corrected <i>FUT2*01.06</i> , c.481G>A, p.Asp161Asn	<i>FUT2*01.06</i> , c.514G>A, p.Asp172Asn
12	Allele	corrected <i>FUT2*01.07</i> , c.665G>A, p.Arg222His	<i>FUT2*01.07</i> , c.698G>A, p.Arg233His
13	Allele	corrected <i>FUT2*01.08</i> , c.685G>A, p.Val229Met	<i>FUT2*01.08</i> , c.6718G>A, p.Val240Met
14	Allele	corrected <i>FUT2*01.09</i> , c.716G>A, p.Arg239Gln	<i>FUT2*01.09</i> , c.749G>A, p.Arg250Gln
15	Allele	corrected <i>FUT2*01.10</i> , c.747_748insGTG, p.249_250insVal	<i>FUT2*01.10</i> , c.780_781insGTG, p.260_261insVal
16	Allele	corrected <i>FUT2*01.11</i> , c.98A>G, c.101T>G, p.Gln33Arg, p.Ile34Arg	<i>FUT2*01.11</i> , c.131A>G, c.134T>G, p.Gln44Arg, p.Ile45Arg
17	Allele	corrected <i>FUT2*01W.01</i> , c.278C>T, p.Ala93Val	<i>FUT2*01W.01</i> , c.311C>T, p.Ala104Val
18	Allele	corrected <i>FUT2*01W.02.01</i> , c.385A>T, p.Ile129Phe	<i>FUT2*01W.02.01</i> , c.418A>T, p.Ile140Phe
19	Allele	corrected <i>FUT2*01W.02.02</i> , c.385A>T, c.617T>G, p.Ile129Phe, p.Val206Gly	<i>FUT2*01W.02.02</i> , c.418A>T, c.650T>G, p.Ile140Phe, p.Val217Gly
20	Allele	corrected <i>FUT2*01W.02.03</i> , c.385A>T, c.841G>A, p.Ile129Phe, p.Gly281Arg	<i>FUT2*01W.02.03</i> , c.418A>T, c.874G>A, p.Ile140Phe, p.Gly292Arg

Track of changes		from	to
1	Version	v6.0 30-JUN-2022	v6.1 31-MAR-2023
21	Allele corrected	<i>FUT2*01W.03</i> , c.853G>A, p.Ala285Thr	<i>FUT2*01W.03</i> , c.886G>A, p.Ala296Thr
22	Allele corrected	<i>FUT2*01W.04</i> , c.617T>C, p.Val206Ala	<i>FUT2*01W.04</i> , c.650T>C, p.Val217Ala
23	Allele corrected	<i>FUT2*01N.01</i> , c.244G>A, c.385A>T, p.Ala82Thr, p.Ile129Phe	<i>FUT2*01N.01</i> , c.277G>A, c.418A>T, p.Ala93Thr, p.Ile140Phe
24	Allele corrected	<i>FUT2*01N.02</i> , c.428G>A, c.739A>G (after termination?), p.Trp143Ter, p.Gly247Ser (after termination?)	<i>FUT2*01N.02</i> , c.461G>A, removed: c.772A>G, p.Trp154Ter, removed: p.Gly258Ser
25	Allele corrected	<i>FUT2*01N.03</i> , c.569G>A, p.Arg190His	<i>FUT2*01N.03</i> , c.602G>A, p.Arg201His
26	Allele corrected	<i>FUT2*01N.04</i> , c.571C>T, p.Arg191Ter	<i>FUT2*01N.04</i> , c.604C>T, p.Arg202Ter
27	Allele corrected	<i>FUT2*01N.05</i> , c.628C>T, p.Arg210Ter	<i>FUT2*01N.05</i> , c.661C>T, p.Arg221Ter
28	Allele corrected	<i>FUT2*01N.06</i> , c.658C>T, p.Arg220Ter	<i>FUT2*01N.06</i> , c.691C>T, p.Arg231Ter
29	Allele corrected	<i>FUT2*01N.07</i> , c.664C>T, p.Arg222Cys	<i>FUT2*01N.07</i> , c.697C>T, p.Arg233Cys
30	Allele corrected	<i>FUT2*01N.08</i> , c.685_686delGT, p.Val229Glyfs*4	<i>FUT2*01N.08</i> , c.718_719delGT, p.Val240Glyfs*4
31	Allele corrected	<i>FUT2*01N.09</i> , c.688_690delGTC, p.Val230del	<i>FUT2*01N.09</i> , c.721_723delGTC, p.Val241del
32	Allele corrected	<i>FUT2*01N.10</i> , c.400G>A, c.760G>A, p.Val134Ile, p.Asp254Asn	<i>FUT2*01N.10</i> , c.433G>A, c.793G>A, p.Val145Ile, p.Asp265Asn
33	Allele corrected	<i>FUT2*01N.11</i> , c.778delC, p.Pro260Leufs*16	<i>FUT2*01N.11</i> , c.811delC, p.Pro271Leufs*16
34	Allele corrected	<i>FUT2*01N.12</i> , c.849G>A, p.Trp283Ter	<i>FUT2*01N.12</i> , c.882G>A, p.Trp294Ter
35	Allele corrected	<i>FUT2*01N.13</i> , c.868G>A, p.Gly290Arg	<i>FUT2*01N.13</i> , c.901G>A, p.Gly301Arg
36	Allele corrected	<i>FUT2*01N.14</i> , c.950C>T, p.Pro317Leu	<i>FUT2*01N.14</i> , c.983C>T, p.Pro328Leu
37	Allele corrected	<i>FUT2*01N.15</i> , c.302C>T, p.Pro101Leu	<i>FUT2*01N.15</i> , c.335C>T, p.Pro112Leu
38	Allele corrected	<i>FUT2*01N.16</i> , c.960A>G (wrong position?), p.Gly247Ser (wrong position?)	<i>FUT2*01N.16</i> , c.772G>A, p.Gly258Ser
39	Allele corrected	<i>FUT2*01N.17</i> , c.412G>A, p.Gly138Ser	<i>FUT2*01N.17</i> , c.445G>A, p.Gly149Ser
40	Allele corrected	<i>FUT2*01N.18</i> , c.818C>A, p.Thr273Asn	<i>FUT2*01N.18</i> , c.851C>A, p.Thr284Asn
41	Reference added		PMID 20570966
42	End Version	v6.0 30-JUN-2022	v6.1 31-MAR-2023

Track of changes		from	to
1	Version	v5.2 18 th April 2019	v6.0 30-JUN-2022
2	Author	created Franz Wagner, April 2019	Franz Wagner, Yanli Ji, June 2022
3	Reviewer	reviewed n.a.	Franz Wagner, C. Gassner, June 2022
4	General	Word version	First Excel map version. Spread-sheets "Intro", "Allele Table", "References", and "Versioning" created.
5	References	moved	Moved reference's texts to tab-sheet 'References'
6	References	numbering	renumbered Abstract (47) to (5)
7	Comment		<p>General comment by Franz Wagner: Since most <i>FUT1</i> alleles are found in trans to a functional <i>FUT2</i> and usually, no expression studies are done, it is extremely difficult to discriminate between <i>FUT1*01W</i> and <i>FUT1*01N</i>.</p> <p>Obviously, the new alleles were assigned in an "optimistic" fashion, i.e. if no data showing a residual activity are present, the allele was assigned to the <i>FUT*01N</i> series although the inactivity of the allele isn't proven. Some of the old <i>FUT1*01W</i> alleles were assigned in a "pessimistic" way, i.e. if no data showing residual activity is absent and the mutation did not destroy protein expression, the allele was assigned to the <i>FUT1*01W</i> series although data showing residual activity are lacking. This change might lead to inconsistencies in the future, but admittedly I have no better solution (as you know, I would have supported a single series for weak or non-functional alleles, but such solution is not in line with the ISBT approach).</p>
8	Allele	changed <i>FUT1*02</i>	changed to <i>FUT1*01.02</i>

Track of changes		from	to
1	Version	v5.2 18th April 2019	v6.0 30-JUN-2022
9	Allele changed	<i>FUT1*02W.01</i>	changed to <i>FUT1*01W.31</i>
10	Allele changed	<i>FUT1*02W.02</i>	changed to <i>FUT1*01W.32</i>
11	Allele changed	<i>FUT1*02W.03</i>	changed to <i>FUT1*01W.33</i>
12	Allele changed	<i>FUT1*02W.04</i>	changed to <i>FUT1*01W.34</i>
13	Allele changed	<i>FUT1*02W.05</i>	changed to <i>FUT1*01W.35</i>
14	Allele changed	<i>FUT1*02N.01</i>	changed to <i>FUT1*01N.26</i>
15	Allele changed	<i>FUT2*02</i>	changed to <i>FUT2*01.02</i>
16	Allele changed	<i>FUT2*03.01</i>	changed to <i>FUT2*01.03.01</i>
17	Allele changed	<i>FUT2*03.02</i>	changed to <i>FUT2*01.03.02</i>
18	Allele changed	<i>FUT2*03.03</i>	changed to <i>FUT2*01.03.03</i>
19	Allele changed	<i>FUT2*04</i>	changed to <i>FUT2*01.04</i>
20	Allele changed	<i>FUT2*05</i>	changed to <i>FUT2*01.05</i>
21	Allele changed	<i>FUT2*06</i>	changed to <i>FUT2*01.06</i>
22	Allele changed	<i>FUT2*07</i>	changed to <i>FUT2*01.07</i>
23	Allele changed	<i>FUT2*08</i>	changed to <i>FUT2*01.08</i>
24	Allele changed	<i>FUT2*09</i>	changed to <i>FUT2*01.09</i>
25	Allele changed	<i>FUT2*10</i>	changed to <i>FUT2*01.10</i>
26	Allele PMIDs added		PMIDs added
27	References added		added authors
28	Allele removed	<i>FUT1*01W.25</i>	obsolete now (provisionally)
29	Allele avoided		The allele name <i>FUT1*01W.30</i> is avoided because of the use of this name in a publication, although this allele had not yet been officially ratified by the ISBT WP.
30	Allele added		<i>FUT1*01W.31</i>
31	Allele added		<i>FUT1*01W.32</i>
32	Allele added		<i>FUT1*01W.33</i>
33	Allele added		<i>FUT1*01W.34</i>
34	Allele added		<i>FUT1*01W.35</i>
35	Allele added		<i>FUT1*01W.36</i>
36	Allele added		<i>FUT1*01W.37</i>

Track of changes		from	to
1	Version	v5.2 18th April 2019	v6.0 30-JUN-2022
37	Allele added		<i>FUT1*01W.38</i>
38	Allele added		<i>FUT1*01W.39</i> (provisionally)
39	Allele changed		<i>FUT1*01N.06</i> added Phenotype H+w and footnote
40	Allele changed		<i>FUT1*01N.16</i> added c.35C>T, PMID 9031499
41	Allele changed		<i>FUT1*01N.20</i> added PMID 34967725
42	Allele changed		<i>FUT1*01N.24</i>
43	Allele changed		<i>FUT1*01N.25</i>
44	Allele changed	<i>FUT1*02N.01</i>	<i>FUT1*01N.26</i> added c.35C>T and rs2071699
45	Allele added		<i>FUT1*01N.27</i> (provisionally)
46	Allele added		<i>FUT1*01N.28</i>
47	Allele added		<i>FUT1*01N.29</i>
48	Allele added		<i>FUT1*01N.30</i>
49	Allele added		<i>FUT1*01N.31</i>
50	Allele added		<i>FUT1*01N.32</i>
51	Allele added		<i>FUT1*01N.33</i>
52	Allele added		<i>FUT1*01N.34</i>
53	Allele added		<i>FUT1*01N.35</i>
54	Allele added		<i>FUT1*01N.36</i>
55	Allele added		<i>FUT1*01N.37</i>
56	Allele added		<i>FUT1*0N.01</i>
57	Allele changed	<i>FUT2*02</i>	<i>FUT2*01.02</i>
58	Allele changed	<i>FUT2*03.01</i>	<i>FUT2*01.03.01</i>
59	Allele changed	<i>FUT2*03.02</i>	<i>FUT2*01.03.02</i>
60	Allele changed	<i>FUT2*03.03</i>	<i>FUT2*01.03.03</i>
61	Allele changed	<i>FUT2*04</i>	<i>FUT2*01.04</i>
62	Allele changed	<i>FUT2*05</i>	<i>FUT2*01.05</i>
63	Allele changed	<i>FUT2*06</i>	<i>FUT2*01.06</i>
64	Allele changed	<i>FUT2*07</i>	<i>FUT2*01.07</i>
65	Allele changed	<i>FUT2*08</i>	<i>FUT2*01.08</i>

Track of changes		from	to
1	Version	v5.2 18th April 2019	v6.0 30-JUN-2022
66	Allele changed	<i>FUT2*09</i>	<i>FUT2*01.09</i>
67	Allele changed	<i>FUT2*10</i>	<i>FUT2*01.10</i>
68	Allele added		<i>FUT2*01.11</i>
69	Allele added		<i>FUT2*01W.04</i>
70	Allele changed		<i>FUT2*01N.04</i> added PMID 11606829
71	Allele added		<i>FUT2*0N.05</i>
72	Allele changed	<i>FUT1*02W.01</i>	<i>FUT1*01W.31</i>
73	Allele changed	<i>FUT1*02W.02</i>	<i>FUT1*01W.32</i>
74	Allele changed	<i>FUT1*02W.03</i>	<i>FUT1*01W.33</i>
75	Allele changed	<i>FUT1*02W.04</i>	deleted, is now <i>FUT1*01N.16</i>
76	Allele changed	<i>FUT1*02W.05</i>	<i>FUT1*01W.34</i>
77	Allele changed		added rs-numbers
78	End Version	v5.2 18th April 2019	v6.0 30-JUN-2022

Track of changes		from	to
1	Version	v5.1 170221	v5.2 18th April 2019
2	Author created	Franz Wagner, February 2017	Franz Wagner, April 2019
3	Reviewer reviewed	n.a.	n.a.
4	General	Word version	Word version
5	Allele Table	allele added	<i>FUT1*01N.21</i>
6	Allele Table	allele added	<i>FUT1*01N.22</i>
7	Allele Table	allele added	<i>FUT1*01N.23</i>
8	Allele Table	allele added	<i>FUT1*01N.24</i>
9	End Version	v5.1 170221	v5.2 18th April 2019